



Global Journal of Scientific Researches

Available online at gjsr.blue-ap.org

©2014 GJSR Journal. Vol. 2(4), pp. 94-97, 30 April, 2014

E-ISSN: 2311-732X

Comparative Study on the Effect of Chloroxylenol and Sodium Chloride on *Brucella* and *Acinetobacter* Growth

Enas I. Jasim

Department of Biology/College of Science /University of Al-Mustansiriya /Iraq

Corresponding Author: Enas I. Jasim

Received: 25 March, 2014

Accepted: 10 April, 2014

Published: 30 April, 2014

ABSTRACT

A study was carried out to investigate the effect of chloroxylenol and sodium chloride on the growth of *Brucella spp.* and *Acinetobacter baumannii* isolates. Results revealed that chloroxylenol was more effect on growth of *Acinetobacter sp.* As compared with *Brucella spp.* The inhibition of the *Acinetobacter* growth was detected at all concentrations (3, 30, 300, 3000, 30000, and 40000)Mg/ ml. Although, *Brucella* isolates resist at 3 and 30 Mg/ ml of chloroxylenol, high growth inhibition was seen at 300 Mg/ ml of chloroxylenol. On the other hand, no difference was found in the effect of chloroxylenol on *Brucella abortus* and *Brucella melitensis*. The results confirmed that the *Brucella spp.* seems to be more susceptible to negative effect of high concentration of NaCl than *A.baumannii*.

Keywords: *Acinetobacter*, *Brucella*, Sodium chloride, Chloroxelenol.

©2014 GJSR Journal All rights reserved.

INTRODUCTION

Infections disease account for about half of the death in tropical countries (Khosravi and Behzadi, 2006). Besides, incidents of epidemic due to drug resistant microorganisms pose enormous public health concerns (Burt and Reinders, 2003). Within the genus *Brucella* there are closely related species that have recognized from many years, this important group causes brucellosis in domestic animals and humans, infection by localized in reticuloendothelial tissues, reproductive organs and less frequently bone and joints (AL-Rodhan, 2005). Microbial resistance to antibiotics especially among *Brucella* species is a major threat to public health (Lang, 1995), mainly in some developing countries like Iraq (Rhaymah et al., 2010). On the other hand *Acinetobacter spp.* is apleomorphic aerobic gram negative bacillus that is widely distributed in nature and causes a wide spectrum of nosocomial infections (Hrenovic and Ivankovic, 2009). It has a capacity for long term survival (up to several months) on most environmental dry surfaces and it has been identified in environmental sampling at military treatment facilities along the Tigris River in Iraq and Kuwait city. Infections in soldiers, shows an increased prevalence of multidrug resistant of *Acinetobacter spp.* in Iraq (Davis, 2005).

Antiseptics and disinfectants are used extensively in hospitals and other health care setting for a variety of hard surface applications, they are on essential part of infection control practices and aid in the prevention of nosocomial infections (Hendry, 2009). Most important factors affecting the survival of enteric bacteria are salinity, nutrient availability, microbial antagonism and antibiotic substances, and bacteria which can tolerate the high salt concentrations up to approximately 10% of sodium chloride are called osmotolerant (Nester, 2004).

This study aimed to investigate the antimicrobial effect of chloroxylenol solution (different concentrations) and compares its effects on *Brucella spp.* and *Acinetobacter spp.* isolates. And the survival of two heterotrophic bacteria in conditions of high salt concentration.

MATERIALS AND METHODS

1. Bacterial strains and cultuer media :

Brucella abortus , *Brucella melitensis* and *Acinetobacter baumannii* isolates were obtained from clinical specimens with brucellosis infections and urinary tract , surgical wound infections or non surgical wound infections in Baghdad hospitals (table-1). The isolates were identified biochemically (biotype analysis) in addition to API-20E analysis (Atlas, 1995) . The isolates were streaked on trypticase soy agar from Himedia (India). For short time preservative at 4 C° and store in 20% glycerol in Brain heart infusion broth (Himedia – India) , - 20C° for long time preservative (10).

2- Study the inhibition of bacterial growth by Chloroxylenol compound.

2-1. Preparation of inoculums .

The isolates subcultured on Brain heart infusion agar (Himedia – India) , for 18- 24 hrs at 37C° , culture purity was checked according to the procedure mentioned by (Forbes et al., 2002) . Ahomogenous bacterial cell suspension (2×10^2) CFU \ml was prepared with turbidity equal to 0.5 Mcfarland (Baron, 1994).

Table 1. The prevalence of *Acinetobacter baumannii* and *Brucella* spp. isolates identified in different clinical cases

Type of isolated bacteria	No. of isolates	Type of sample
<i>Brucella melitensis</i>	2	Blood from brucellosis patients
<i>Brucella abortus</i>	3	Blood from brucellosis patients
<i>Acinetobacter baumannii</i>	5	Wounds infection
<i>Acinetobacter baumannii</i>	5	Urine from urinary tract infection patients
Total No. of isolates	15	

2-2. Chloroxylenol solution

Chloroxylenol is the principal ingredient in Dettol. It prepared by diluting with distilled water , solution of (0.2)% Chloroxylenol of commercial grade , adding to distilled water and adjusting to the final strength of Dettol to (3 , 30 , 300 , 3000 , 30.000 and 40.000)Mg\ml.

2 -3. Minimal inhibitory concentration.

The effect of Chloroxylenol dilutions on the growth of *Brucella* and *Acinetobacter* isolates was determined by a macrobroth dilution method (Messenger et al., 2001).Each tube contained 1ml of 6 concentration of Chloroxylenol (5:1) as doubling dilution in the range (3 – 40.000)Mg \ml .

The tubes were inoculated with 100 *ml* of an 18- 24 hrs culture diluted to yield a final inoculum of 10^8 CFU \ml. Brain heart infusion broth was used both as diluent and as culture medium . The MICs value Mg \ml was recorded as the lowest concentration in which no visible growth was apparent after incubation for 24 hrs at 37C°.

3 - Inhibition of bacterial growth by sodium chloride.

3 – 1. Preparation of inoculum was done as in (Burt and Reinders, 2003; Khosravi and Behzadi, 2006).

3 – 2. Experimental procedures as in (Hrenovic and Ivankovic, 2009) .

Nutrient broth obtained from himedia – Indin was used and analytical sodium chlorid (BDH – England) were added in the flasks to obtain the concentration of (1, 2 , 3 , 4 , 5 , 6 , 7 , 8 , 9 , 10 , 11) % of NaCl in nutrient broth . the pH value of broth media was adjusted to 7.0 -⁺0.1 using pH – meter .

One ml of resuspended biomass was inoculated into Erlenmeyer flasks with ranging concentrations of NaCl in 100 ml of nutrient broth. The flasks were sealed and incubated for 24 hrs in 37C° . starting number of viable count was determined before incubation. And the number of viable cells in each bottle was determined after 24 hrs . of incubation . as colony forming units . A 1 ml of bacterial suspension was aseptically taken from each bottle and serially diluted (10^{-1} , 10^{-2} , 10^{-3}). Volumes of 0.1 ml were then aseptically inoculated on to nutrient agar plates (spread –plate method) . After the incubation 24 hrs . The bacterial colonies were counted and the number of cells was reported as CFU/ml . All measurements were done in triplicate and mean values are presented.

RESULTS AND DISCUSSION

The results shows that the antibacterial activity of the chloroxylenol product varies with both organisms (*Brucella* spp. and *Acinetobacter* sp.) . This variation is particularly observable against *Acinetobacter baumannii* and to a lower extent against *Brucella* spp. while virtually no variation is appear with *B. melitensis* and *B.abortus*. The MIC of the chloroxylenol against these organisms are listed in Table (2). Chloroxylenol range between (3-40000) Mg/ml for *A.baumannii* , while it range between 30

to 300 Mg/ml for *B.abortus* and *B.melitensis* , since preliminary studies usually show MIC of ≤ 1 Mg/ml (Hassain, 2011) , but against streptococcus mutant but not against this study organisms .

Chloroxylenol as phenolic compound can inhibit the Adenosine triphosphatase (ATPase) which is an important enzyme that is linked to cytoplasmic membrane and thus can inhibit the procese of returning potassium ions into cells in exchange for sodium and hydrogen ions , (Autio, 2008) . also inhibits metabolic enzymes such as phosphoenol pyruvate phosphotransferase , which lead to inhibit and kill the organisms (Axmann and Brex, 2008).

Table 2. In vitro activity of chloroxylenol as different concentration against *Brucella* spp. and *Acinetobacter baumannii* mention as MIC.

Type of isolates	3	30	300	3000	30000	40000
<i>B. melitensis</i>	+	-	-	-	-	-
<i>B. Abortus</i>	+	-	-	-	-	-
<i>A. baumannii</i>	-	-	-	-	-	-

- : No growth
+ : growth

In the second set of experiments nutrient rich medium was used for bacterial growth . In the control reactors without addition of Nacl , both bacteria (*Brucella* and *Acinerobacter*) were intensively multiplying and the number of both bacteria after 24 hrs increased for one order of magnitude when compared to starting number (3.75×10^2 CFU/ml) table – 3 . The bacteria most likely similar growth trend was obtained in reactors containing (1 ,2 ,3 ,4 ,5) % of Nacl .

While at (6, 7, 8 , 9 and 10) % of Nacl the number of *A.baumannii* was lowed when compared to starting number and decreased further in 10 % with no growth at 11% of Nacl . Indicating decay of cells.

For *Brucella* spp. the growth at different concentration of Nacl was similar to that obtained for *A. baumannii*. Concerning, *Brucella* spp. there was no growth at (8 , 9 , 10) % of Nacl . Based on the results, it seems that *Brucella* spp. was more susceptible to negative effect of high concentration of Nacl than *A.baumannii* . Increased concentration of Nacl caused a failure in bacterial multiplication and decay of bacterial cell and death of all cells in the growth media . These finding agree with Stein, (2000).

Table 3. The effect of sodium chloride on the growth of *Brucella* spp. and *Acinetobacter baumannii*

Type of isolates	No. isolates	Concentration of Nacl %											Start No.
		CFU/ ml $\times 10^2$											
		1	2	3	4	5	6	7	8	9	10	11	
<i>A.baumannii</i>	10	3.60	3.59	3.50	3.35	3.35	3.00	2.30	1.60	1.10	0.8	0	3.75
<i>B.abortus</i>	3	3.21	2.48	1.77	1.06	0.35	0.7	0	0	0	0	0	3.26
<i>B. melitensis</i>	2	3.50	3.28	3.15	2.13	0.71	0.15	0.2	0	0	0	0	3.58

REFERENCES

AL-Rodhan M A .2005 . Survey of Brucellosis in cattle in AL-Diwaniya City . Iragi J, Vet . Sci , 4 (2) : 13-17.

Atlas RM, Park L and Brown A .1995 . Laboratory Manual of Experimental Microbiology . 1 ed . Mosby , Co. USA.

Autio J. 2008. The role of chlorohexidine in caries prevention . Oper. Dant , 33 (6) : 710-716 .

Axmann D and Brex M. 2008 . Effect of of xylitol / chlorohexidine versus xylitol as single rinses on intial biofilm formation of streptococci . Quintessence Int ., 39(1) : 7-22 .

Baron EJ, Finegold SM and Peterson I.LR. 1994. Baily and Scott's Diagnostic Microbiology. 9th-ed. Mosby , Co.USA .

Burt AS and Reinders RD. 2003. Antibacterial activity of Selected plant essential oils against *Escherichia coli* O157 :H7 . Lett .Appl . Microbial , 36 : 162-167.

Davis KA, Moran KA, McAllister CK and Gray PJ .2005. Multidrug – Resistant *Acinetobacter* Extremity Infections in Soldiers, Emerg. Infect. Dis , 11 : 1218- 1224.

Forbes BA, Saham DF and Weissfeld AS. 2002. Baily and Scott's Diagnostic Microbiology . 11thed. Mosby, Inc. St. Louis . USA.

Hassain NS. 2011. Study metabolic activities of Mutans streptococci isolated from the mouth of diabetes mellitus type 1 patients compared to the normal individuals , thesis . MSc. college of dentistry –Baghdad university .

Hendry ER, Worthington J, Conway BR and Lambert PA. 2009. Antimicrobial efficacy of eucalyptus oil and 1,8-cineol alone and in combination with chlorhexidine digluconate against microorganisms grown in plank tonic and biofilm cultures . J . Antimicro . chemo , 64: 1219-1225.

Hrenovic J and Ivankovic T. 2009 .Survival of *Escherichia coli* and *Acinetobacter junii* at various concentrations of sodium chloride. Eur . Asia J . Bio . Sci , 3 (18) : 144-151 .

Khosravi A and Behzadi A . 2006 . Evaluation of the antimicrobial activity of the seed hull of *Quercus baranti* on some gram negative bacteria . Pak . J . Med . Sci , 22 : 429-432 .

Lang RM, Banai MU and Rubinstein E . 1995. Brucellosis . Antimicrob . Agen , 5 (2) : 205-208.

- Messenger S, Goddard PA and Dettmar PW. 2001. Determination of the antibacterial activity of several antiseptics tested on skin by an ex-vivo test. *J. Med. Microbiol*, 50 :284-292.
- Nester EW, Anderson DG, Roberts CE and Nester MT .2004. *Microbiology : a human perspective* . McGraw. Hill. Inc, New York.
- Rhaymah MS, AL-Saad AK and AL-Hankawe KH. 2010. Diagnosis of bovine brucellosis in Mosul city by indirect ELISA and conventional serological test . *Iraqi J. Vet . Sci* , 24 (1) : 1-6 .
- Stein R . 2000 . *Know the enemy : Under standing the basis of food borne pathogens* . *Meat Poultry* , 10 : 46-51 .