



## Global Journal of Scientific Researches

Available online at [gjsr.blue-ap.org](http://gjsr.blue-ap.org)

©2019 GJSR Journal. Vol. 7(2), pp. 23-26, 19 October, 2019

E-ISSN: 2311-732X

# Study of Skin Receptor of Cattle through Histological Aspect

Jamil Oglu<sup>1</sup>, Forough Talaeizadeh<sup>2</sup> and Ahmad Riahi<sup>3\*</sup>

1- Agricultural Research, University of Baku, Azerbaijan

2- Department of Educational Science and Psychology, Islamic Azad University Marvdasht Branch, Iran

3- Department of Biology, Payam Noor University, Tehran, Iran

**Corresponding Author:** Ahmad Riahi

*Received: 20 September, 2019*

*Accepted: 01 October, 2019*

*Published: 19 October, 2019*

---

### ABSTRACT

Sarabi cattle are found in Iran and Azerbaijan. They are used for both dairy and meat production. A histological study was carried out on the teat of Sarabi cattle for the first time in Iran. A total number of five adult and healthy cows were selected and after slaughtering, tissue samples were taken from different parts of the teat skin. After fixation in 10% formalin, they were transferred into the tissue processor. Paraffin blocks were made and thin sections of 6 $\mu$ , were cut. The sections were subjected to stain by Hematoxylin and Eosin. They were studied under the light microscope and photo micrographs were taken. The results show that the teat consists of teat sinus, teat wall, teat canal and teat orifice. Teat wall includes three separable strata: skin, middle layer and mucosal layer. Teat skin contains an epidermis and dermis, but there is no evidence of hypodermis existence in contrast with the other parts of the body. Dermis is formed from networks of collagen fibers, blood vessels, circular smooth muscle and nerve fibers. Meissner's corpuscles are found in the dermal papillae particularly in the connective tissue of teat canal. Myelinated nerve fibers are also seen in this region.

**Keywords:** *Histology, Sarabi cattle, Sensory receptors, Skin, Teat.*

©2019 GJSR Journal All rights reserved.

---

### INTRODUCTION

The Sarabi indigenous cow is one of the major regional dual purpose cows in Iran, which has a strong adaption to cold weather, diseases and parasites. The breed was imported from Gharabagh and breeds in Sarab, Mianeh and Ardebil regions of Iran (Khansefid, 2009). Individuals of this breed are typically red (Mason, 1996). Although Sarabi is a dual purpose cow, the milk production of this breed is noticeably higher than other Iranian vernacular cow breeds (Khansefid, 2009). The average milk yield in this breed is 10 kilograms daily with an approximately 4.5% fat. Also production up to 20 kilograms daily can be seen individually (Mason, 1996).

Milk let-down is the natural process used by the cow to help remove milk from the udder. This process is brought about by the release of the oxytocin hormone from posterior pituitary gland under control of hypothalamus, following a suitable stimulus being received by the cow. The stimulus may be visual, heard or felt and should be predictable and consistent at every milking. Most importantly it should not induce fear in cows. It is commonly thought that genetic selection has ensured that most commercial cows will let-down freely without physical contact with the teats. Handling of the teats however, is a strong stimulus for let-down with research showing that at least 15 seconds of massage per cow is required to effect an appreciable change in milking characteristics. After release, oxytocin travels through the blood stream and has a direct effect on small muscle cells that surround the milk-producing cells in the udder. Oxytocin causes these myoepithelial cells to contract and squeeze the milk into the milk ducts and so towards the teats. The pressure of the milk being forced into the ducts and down towards the teat causes the teat to swell with milk and become "plump" with milk. It takes between 60 and 90 seconds for the teats to become plump with milk after let-down has been initiated (Klindworth, 2003). A close study on the sensory receptors of teat skin would be very important to understand the physiology and pathology of milk let-down. Histological study of sensory receptors of the teat skin in Sarabi cattle was carried out for the first time in Iran.

## 2. Methodology

In order to histological study of the sensory receptors in areola and skin of the teat of Sarabi cattle, five adult and healthy cows were selected and after slaughtering, tissue samples were taken from different parts of the skin and areola of the teat. After fixation in 10% formalin, they were transferred into the tissue processor. Paraffin blocks were made and thin sections of 6  $\mu$ , were cut. The sections were subjected to stain by Hematoxylin and Eosin. They were studied under the light microscope and photo micrographs were taken.

## 3. Results

### 3.1 Macroscopic investigation

Each quarter has a separate teat with 7–8 cm length and 3 cm diameter. The teats vary in shape from cylindrical to conical. The results show that the teat consists of teat sinus, teat wall, teat canal and teat orifice. Each teat has a strong structure that protects it from excessive pressure due to the calf sucking, milking machine or hand milking. Despite the skin of the mammals, teat skin in Sarabi cattle is tightly attached to the underlied tissues and thus promotes teat resistance.

### 3.2 Microscopic investigation

Teat wall includes three separable strata: skin, middle layer and mucosal layer. Teat skin contains an epidermis and dermis, but there is no evidence of hypodermis existence in contrast with the other parts of the body. Epidermis consists of keratinized stratified squamous epithelium and the thickness increases towards the teat orifice. Dermis is formed from networks of collagen fibers, blood vessels, circular smooth muscle and nerve fibers. Dermal papillae projects into the epithelial undulations at the dermo-epidermal junction. Meissner's corpuscles are found in the dermal papillae particularly in the connective tissue of teat canal (Figure 1).




Figure 1. Teat dermis (H&E  $\times$ 100).  
Arrow: Blood vessels.




Figure 2. Dermal papillae (H&E  $\times$ 400).  
Arrow: Meissner's corpuscles.

Myelinated nerve fibers are also seen in this region (Figure 2). Middle layer (Second layer) is the thickest layer of the teat wall that contains numerous smooth muscle bundles, developed vascular plexus and peripheral nerve fibers (Figure 3).




Figure 3. Middle layer (H&E  $\times$ 400).  
Arrow: Peripheral nerves.

Mucosal layer is the innermost layer of the teat wall. Teat canal epithelium was thicker than teat skin epithelium (150 vs. 110  $\mu$ ) with a remarkable increase in keratin and stratum granulosum.

#### 4. Discussion and Conclusion

The bovine teat canal is highly specialized in its unique function of preventing both leakage of milk and entry of bacteria and thereby plays a major role in the defence of the udder against mastitis (Paulrud, 2005). The surface of the bovine and porcine teat is covered by glabrous, aglandular, stratified squamous epithelium (Frandsen et al., 2009). In the mare, bitch and queen abundant sebaceous glands and fine hairs are present (Aughey, 2001). Nerve ends are located in the teat, in such a way that responds well to the hormonal neural stimuli. Characteristic of teat receptors in rabbit were studied by recording electrophysiologic wave techniques for afferent nerves and it was shown that teats are provided with mechanical receptors sensitive to suckling and pressure (Ramakrishna, 2004). Hebb and Linzell (1970) reported a great number of nerve fibers in the arteries and arteriols that forms a plexus in the vessel wall. Some of these fibers exist near the teat canal wall, but none of them ends to the alveolus and myoepithelial cells. In addition to the nerve fibers that end to the ducts, it was found many nerve fibers near the teat orifices. It was shown that goats can differentiate left and right udder stimulation. Numerous sensory nerve fibers in the middle layer conduct impulses to the CNS and oxytocin is released from neurohypophysis. It was indicated that there are mechanical receptors in the teat of rabbits (Hurley, 1989; Turner, 1996). In the distal portion of the teat, there is a high density of circular smooth muscle surrounding the teat canal. It was indicated that the aggregation of muscular tissue in this area in heifers is less than the delivered cows (Linzell, 1971; William, 1993).

#### REFERENCES

1. Aughey E, Frye FL, 2001. Comparative Veterinary Histology. Manson Publishing Ltd. London. UK.
2. Frandsen RW, Wilke WL, Dee Fails A, 2009. Anatomy and physiology of farm animals. Wiley Blackwell.
3. Hebb CA, Linzell LJ, 1970. Innervation of the mammary gland. A Histochemical study in the rabbit. *Histochem J.* 2(6): 491-505.
4. Hurley WL, 1989. Mammary gland Function during involution. *J Dairy Sci.* 72(6): 1637-1646.
5. Khansefid SM, 2009. An introduction to Sarabi cow breed – a vernacular cow breed of Iran. *Nolan Special Magazine on Dairy cows.* No 3/Jan-Mar.
6. Klindworth, D., Campbell, J. and Greenall, R. (2003) CowTime Guidelines for milk harvesting. Dairy Research and Development Corporation. Available on the Cow Time Website, [www.cowtime.com.au](http://www.cowtime.com.au).
7. Linzell LJ, Peaker M, 1971. *Physiol Rev.* 51(3): 564-597.
8. Mason IL, 1996. A World Dictionary of Livestock Breeds, Types and Varieties. 4<sup>th</sup> edn. CAB International.
9. Paulrud CO, 2005. Basic Concepts of the Bovine Teat Canal. *Vet Res Commun.* 29(3): 215-245.
10. Ramakrishna V, Gadre KM, 2004. A Systemic Histology of Domestic Animals, 2<sup>th</sup> edn. Jaypee Brothers Medical, New Delhi.
11. Turner CE, 1996. The Anatomy of the Udder of Cattle and Domestic Animals. *Biol Reprod.* 55: 671- 675.
12. William GL, McVey WR, Hunter JF, 1993. Mammary somatosensory pathways are not required for suckling-mediated inhibition of luteinizing hormone secretion and delay of ovulation in cows. *Biol Reprod.* 49: 1328-1337.
13. Aida Badamchi Shabestari, B. A. A., Maryam Shekarchi, Seyed Mojtaba Mostafavi. (2018). Development of Environmental Analysis for Determination of Total Mercury in Fish Oil Pearls by Microwave Closed Vessels Digestion Coupled with ICP-OES. *Ekoloji*, 27(106), 1935.
14. Aida Badamchi Shabestari, S. M. M., Hanieh Malekzadeh. (2019). Force Degradation Comparative Study on Biosimilar Adalimumab and Humira. *Revista Latinoamericana de Hipertensión*, 13(06), 496-509.
15. Al Jamal, A., & Al Yousef, M. (2018). Phytochemical Analysis of Some Herbal Medicines. *MedBioTech Journal*, 02(03), 82-84. doi:10.22034/MBT.2018.76928
16. Amanlou, M., & Mostafavi, S. M. (2017). In silico screening to aim computational efficient inhibitors of caspase-9 by ligand-based pharmacophore modeling. *MedBioTech Journal*, 01(01), 34-41. doi:10.22034/mbt.2017.60337
17. Amir Yaghoubi Nezhad, S. H., Atefeh Mehrabi Far, Masoumeh Piryaee, Seyed Mojtaba Mostafavi. (2019). Investigation of Shigella Lipopolysaccharides Effects on Immunity Stimulation of Host Cells. *International Transaction Journal of Engineering, Management, Applied Sciences and Technologies*, 10, 465.
18. Gowda, S., & Kumar, A. (2018). Measurement the Antifertility Potential in Male Rats Exposed to The Electromagnetic Signals Emitted by Cellular Mobile Telephony. *MedBioTech Journal*, 02(03), 108-112. doi:10.22034/MBT.2018.76938
19. Grizni, A. (2019). Synthesis, Determination of the Structure and Electrochemical Performance of the Palladium-Cobalt Alloy Catalyst Based on the Graphene for usage in the direct alcohol fuel cells with ethanol fuel. *MedBioTech Journal*, 03(01), 22-25. doi:10.22034/mbt.2019.80837
20. Gupta, P., & Kumar, R. (2018). PCR Analysis of a Chinese Herbal in Some Indian States. *MedBioTech Journal*, 02(02), 47-49. doi:10.22034/MBT.2018.76912
21. Heidari, S., Imani, M., & Mostafavi, S. M. (2017). A Validated and Rapid HPLC Method for Quantification of Human Serum Albumin in Interferon beta-1a Biopharmaceutical Formulation. *MedBioTech Journal*, 1(01), 29.
22. Heidary, S., & Riahi, A. (2018). Study of  $\alpha$ -1 Antitrypsin Serum and its Effects on Chronic Inflammation in Diabete Patients. *MedBioTech Journal*, 02(01), 01-07. doi:10.22034/MBT.2018.60861

23. Jafari, S., & Mostafavi, S. A. (2019). Investigation of nitrogen contamination of important subterranean water in the plain. *MedBioTech Journal*, 03(01), 10-12. doi:10.22034/mbt.2019.80826
24. Jamal, G., & Rahman, S. (2018). Marker-Analysis of Leaf Resistance Bacterial in Rice. *MedBioTech Journal*, 02(03), 87-94. doi:10.22034/MBT.2018.76931
25. Jamal, M., & Oglu, A. (2018). Review of The Basis for Oncolytic Virotherapy and Development of the Genetically Modified Tumor-Specific Viruses. *MedBioTech Journal*, 02(03), 95-102. doi:10.22034/MBT.2018.76932
26. Lebanov, P., & Nesterenko, V. (2017). Study of Physicochemical Characters of in-situ Gel of Tramadole Hydrochloride. *MedBioTech Journal*, 01(02), 52-59. doi:10.22034/MBT.2017.60368
27. Mehrabifar, A., & Rahmati, M. (2017). Significant Increasing of Gastrointestinal Dissolution of Aceclofenac via Florite. *MedBioTech Journal*, 01(01), 9-14. doi:10.22034/MBT.2017.60328
28. Mostafavi, S. M. (2015). 3D Graphene Biocatalysts for Development of Enzymatic Biofuel Cells: A Short Review. *Journal of NanoAnalysis*, 2(2), 57-62.
29. Mostafavi, S. M., Bagherzadeh, K., & Amanlou, M. (2017). A new attempt to introduce efficient inhibitors for Caspas-9 according to structure-based Pharmacophore Screening strategy and Molecular Dynamics Simulations. *MedBioTech Journal*, 01(01), 1-8. doi:10.22034/mbt.2017.60325
30. Mostafavi, S. M., Malekzadeh, H., & Taskhiri, M. S. (2019). In Silico Prediction of Gas Chromatographic Retention Time of Some Organic Compounds on the Modified Carbon Nanotube Capillary Column. *J. Comput. Theor. Nanosci*, 16, 151-156.
31. Ologun, M., & Salman, Y. (2017). Quantitative Determination of Pefloxacin Mesylate through UV Method and Study of Gelling Form of Pefloxacin. *MedBioTech Journal*, 01(02), 79-84. doi:10.22034/MBT.2017.60378
32. Parvianian, S., Mostafavi, S. M., & Aghashiri, M. (2016). Multifunctional Nanoparticle Developments in Cancer Diagnosis and Treatment. *Sensing and Bio-Sensing Research*, 1(2), 22.
33. Pasban, A., Mostafavi, S. M., Malekzadeh, H., & Mohammad Nazari, B. (2017). Quantitative Determination of LPG Hydrocarbons by Modified Packed Column Adsorbent of Gas Chromatography Via Full Factorial Design. *Journal of NanoAnalysis*, 4(1), 31-40.
34. Putra, A., & Riahi, S. (2018). The Simultaneous Effects of some Herbal Mixtures on Methimazole Medicine for Thyroide Treatment. *MedBioTech Journal*, 02(01), 08-13. doi:10.22034/MBT.2018.60975
35. Riahi, A. (2018). Oil-in-Water Emulsion-Solvent Evaporation Fabrication of Polycaprolactone Containing Piroxicam for Drug Release. *MedBioTech Journal*, 02(01), 14-20. doi:10.22034/MBT.2018.60876
36. Samira Eissazadeh, M. P., Mohammad Sadegh Taskhiri, & Mostafavi, S. M. (2019). Improvement of Sensitivity of Antigen-Antibody Detection of Semen Through Gold Nanoparticle. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 10(1), 144.
37. Samira Eissazadeh, M. P., & Mostafavi, S. M. (2019). Measurement of Some Amino Acid Using Biosensors Based on Silicon-Based Carbon Nanotubes. *Journal of Computational and Theoretical Nanoscience*, 16, 1.
38. Samira Eissazadeh, S. M. M., Masoumeh Piryaeei, & Taskhiri, M. S. (2019). Application Of Polyaniline Nanostructure Based Biosensor For Glucose And Cholesterol Detection. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 10(1), 150.
39. Seyed Mojtaba Mostafavi, H. M., & Taskhiri, M. S. (2019). In Silico Prediction of Gas Chromatographic Retention Time of Some Organic Compounds on the Modified Carbon Nanotube Capillary Column. *Journal of Computational and Theoretical Nanoscience*, 16(151), 156.
40. Seyed Mojtaba Mostafavi, M. P., Ahmad Rouhollahi, & Mohajeri, A. (2014a). Separation and Quantification of Hydrocarbons of LPG Using Novel MWCNT-Silica Gel Nanocomposite as Packed Column Adsorbent of Gas Chromatography. *Journal of NanoAnalysis*, 1(01), 01.
41. Seyed Mojtaba Mostafavi, M. P., Ahmad Rouhollahi, & Mohajeri, A. (2014b). Separation of Aromatic and Alcoholic Mixtures using Novel MWCNT-Silica Gel Nanocomposite as an Adsorbent in Gas Chromatography. *Journal of NanoAnalysis*, 1(01), 11.
42. Shamsipur, M., Beigi, A. A. M., Teymouri, M., Poursaberi, T., Mostafavi, S. M., Soleimani, P., . . . Tash, S. A. (2012). Biotransformation of methyl tert-butyl ether by human cytochrome P450 2A6. *Biodegradation*, 23(2), 311-318.
43. Sokola, J., & Yonakolo, J. (2018). Study of Nutritional Values of Some Traditional Foods in Nigeria. *MedBioTech Journal*, 02(02), 65-68. doi:10.22034/MBT.2018.76924.
44. Wat, P. (2017). Pb and Cd Detection in some Pharmaceutical Products through Electrochemical Method. *MedBioTech Journal*, 01(01), 22-28. doi:10.22034/MBT.2017.60335.
45. Z. Man, A. G. E., S. M. Mostafavi, & Surendar, A. (2019). Fuel oil characteristics and applications: economic and technological aspects. *Petroleum Science and Technology*.